

Peck, S. B., Herrera, H. W. (2012). CDF Checklist of Galapagos Cockroaches, Mantids and Termites - FCD Lista de especies de Cucarachas, mantidos y termitas de Galápagos. *In*: Bungartz, F., Herrera, H., Jaramillo, P., Tirado, N., Jiménez-Uzcátegui, G., Ruiz, D., Guézou, A. Ziemmeck, F. (eds.). Charles Darwin Foundation Galapagos Species Checklist - Lista de Especies de Galápagos de la Fundación Charles Darwin. Charles Darwin Foundation / Fundación Charles Darwin, Puerto Ayora, Galapagos: <http://checklists.galadarwin.webfactional.com/terrestrial-invertebrates/dictyoptera/> Last updated 20 May 2012.

CDF Checklist of Galapagos Cockroaches, Mantids and Termites

Stewart B. Peck, Henri W. Herrera

Contribs.: Sandra Abedrabbo, Fabián Bersosa, Ruth Boada, Charlotte Causton, Germania Estévez, Lilian Guzmán, Henri W. Herrera, María T. Lasso, María Piedad Lincango, E. G. Linsley, Yale Lubin, Alejandro Míeles, Renato Oquendo, Christine E. Parent, Helmut W. Rogg, Lázaro Roque-Álbelo, Leslie Usinger.

20 May 2012

This checklist is automatically generated using Version 3.0 of the online database CDF Galapagos Species Checklist.

CockroachesCockroaches are archaic, hardy, and successful insects and are most diverse in tropical countries.

They are usually general feeders and many do well in the presence of humans. Seemingly only three or four species naturally colonized the Galapagos on their own, by rafting. Considering the rich Neotropical fauna, this seems a surprisingly low number.

Another 11 species were probably brought by humans, some perhaps as early as the first European landing in 1535. **Mantids**There are eight families of Mantids worldwide. Most families, genera, and species are tropical. All are predators on other insects. While the fauna of tropical America is very rich, there is only one species in Galapagos. They are most often seen at night. Many mantid females are flightless, as is the Galapagos species. **Termites**Termites are social insects which live in colonies in the subtropics and tropics. They feed on dead plant material, especially wood and thus play an important role as detritivores in the regions where they occur.

For marine species distribution data cited in the CDF Galapagos Checklists refer to the five main bioregions of the archipelago (Far Northern, Northern, Western, South Eastern and the Elithabeth Bay Bioregion). For the terrestrial species the more than 120 islands, islets and small rocks have been aggregated into Islands Groups, thus, for example, the island group "Santa Cruz" includes smaller islands like Santa Fé, Plaza Norte, Plaza Sur, Baltra, Daphne Mayor, Daphne Minor, and others.

Please be aware that the distribution data presented here is automatically generated from specimen records and does not always accurately reflect the known distribution for all species.

IUCN red-list assessments presented here may deviate from the global IUCN list for the following reasons:

- for well known species groups like vascular plants or vertebrates updates proposed to the IUCN are shown instead of the outdated, but currently accepted status;
- for poorly known species groups (e.g., lichenized fungi) a general assessment is currently not possible and the list presented here is a regional red-list list for Galapagos archipelago.

Names of taxa included in this checklist: 27 (26 Accepted , 1 Unidentified Taxon).

Origin of the taxa included: 11 Accidental, 5 Questionable Native, 7 Endemic.

1. *Anaplecta lateralis* (Burmeister, 1838)
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Introduced, Accidental.
Galapagos Distribution: Isabela, Santa Cruz, Santiago.
References: Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).
2. *Blaberus parabolicus* Walker, 1868
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Introduced, Accidental.
Galapagos Distribution: San Cristóbal, Santa Cruz.
References: Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).
3. *Blattella germanica* (Linnaeus, 1767)
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Introduced, Accidental.
Galapagos Distribution: Española, Santa Cruz.
References: Hebard, M. et al. (1920), Linsley, E.G. et al. (1966), Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).
4. *Chorisoneura carpenteri* Roth, 1988
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Native, Endemic.
Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santa Fé, Santiago.
References: Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).
5. *Chorisoneura cristobalensis* Roth, 1992
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Native, Endemic.
Galapagos Distribution: San Cristóbal.
References: Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).
6. *Cryptotermes darwini* (Light, 1935)
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Native, Endemic.
Galapagos Distribution: Floreana, Isabela, Santa Cruz, Santiago, Wolf.
References: Bahder B.W. et al. (2009), Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).
7. *Cryptotermes fatalus* (Light, 1935)
Taxon status: Accepted name; taxon occurs in Galapagos.
 Syn.: *Kalotermes occidentalis*
Origin: Introduced, Questionable Native.
Galapagos Distribution: Isabela, Santiago.
References: Bahder B.W. et al. (2009), Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).
8. *Galapagia solitaria* Scudder, 1893
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Native, Endemic.
Galapagos Distribution: Española, Floreana, Isabela, San Cristóbal, Santa Cruz, Santa Fé, Santiago.
References: Hebard, M. et al. (1920), Linsley, E.G. et al. (1966), Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).
9. *Heterotermes convexinotatus* (Snyder, 1924)
Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Heterotermes orthognathus*. Nickle Collins (1992) list this as *Heterotermes aureus convexinotatus* for Panama.

Origin: Introduced, Questionable Native.

Galapagos Distribution: Santa Cruz.

References: Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).

10. *Heterotermes orthognathus* Light, 1933

Taxon status: Accepted name; taxon occurs in Galapagos.

Galapagos Distribution: Unknown.

References: Light, S.F et al. (1935), Linsley, E.G. et al. (1966).

11. *Holocompsa nitidula* (Fabricius, 1781)

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Introduced, Questionable Native.

Galapagos Distribution: Isabela, Santa Cruz.

References: Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).

12. *Holocompsa* sp.

Taxon status: Taxon not identified to species, subspecies, form or variety.

Origin: Introduced, Questionable Native.

Galapagos Distribution: Santa Cruz.

References: Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).

13. *Incisitermes galapagoensis* (Banks 1901)

Taxon status: Accepted name; taxon occurs in Galapagos.

Galapagos Distribution: Unknown.

References: Bahder B.W. et al. (2009).

14. *Incisitermes pacificus* (Banks, 1901)

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Incisitermes galapagoensis*, *I. immigrans*. Peck (1996): "Light (1935) states that this species should be considered as a senior synonym of *I. tabogae* Snyder, 1924, but Nickle Collins 1992 use *I. tabogae* as a valid species. *I. galapagoensis* is a supposed endemic species from Islas Wolf and Genovesa. This species cannot be recognized by the descriptions. Cotypes are in the USNM, but Constantino (1998) states that the holotype adult was in CASC, and was destroyed. The distribution of *I. immigrans* is supposed to be Central Pacific Islands, (Hawaii, Marquesas, Fanning, Jarvis), El Salvador, Panama, and Peru; and widespread in the Galápagos. Light (1935) gives characters of soldiers to separate *I. immigrans* from *I. tabogae* (= *I. pacificus*), but I find both "species" of soldiers in the same colony. I have listed above all island records of *Incisitermes* as *I. pacificus*, since it is the oldest available name for the Galápagos. My specimens are deposited in the USNM, and these may help solve the questions about the correct application of these names."

Origin: Introduced, Questionable Native.

Galapagos Distribution: Fernandina, Floreana, Genovesa, Isabela, Marchena, Pinta, Pinzón, San Cristóbal, Santa Cruz, Santa Fé, Santiago, Wolf.

References: Bahder B.W. et al. (2009), Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).

15. *Ischnoptera peckorum* Roth, 1988

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

Galapagos Distribution: Santa Cruz.

References: Peck, S.B. et al. (1996), Peck, S.B. et al. (1990), Peck, S.B. et al. (2001), Roth, L.M. et al. (1988).

16. *Ischnoptera santacruzensis* Roth, L. M. 1992

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

Galapagos Distribution: Isabela, Santa Cruz.

References: Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).

17. *Ischnoptera snodgrassii* (McNeil, 1901)

Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Native, Endemic.

- Galapagos Distribution:** Fernandina, Floreana, Isabela, Santa Cruz, Santiago.
References: Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (2001).
18. *Kaloterme fatulus* Light, 1935
Taxon status: Accepted name; taxon occurs in Galapagos.
Galapagos Distribution: Unknown.
References: Light, S.F et al. (1935), Linsley, E.G. et al. (1966).
19. *Kaloterme immigrans* Snyder, 1922
Taxon status: Accepted name; taxon occurs in Galapagos.
Galapagos Distribution: Unknown.
References: Light, S.F et al. (1935), Linsley, E.G. et al. (1966).
20. *Nauphoeta cinerea* (Oliver, 1789)
Taxon status: Accepted name; taxon occurs in Galapagos.
 Syn.: *Nauphoeta bivittata*, *N. circumvagans*, *N. laevigata*?
Origin: Introduced, Accidental.
Galapagos Distribution: Floreana, San Cristóbal.
References: Hebard, M. et al. (1920), Linsley, E.G. et al. (1966), Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).
21. *Periplaneta americana* (Linnaeus, 1758)
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Introduced, Accidental.
Galapagos Distribution: Floreana, Genovesa, Isabela, San Cristóbal, Santa Cruz.
References: Linsley, E.G. et al. (1966), Peck, S. et al. (1986), Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).
22. *Periplaneta australasiae* (Fabricius, 1775)
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Introduced, Accidental.
Galapagos Distribution: Floreana, Isabela, San Cristóbal, Santa Cruz.
References: Hebard, M. et al. (1920), Linsley, E.G. et al. (1966), Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).
23. *Periplaneta brunnea* Burmeister, 1838
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Introduced, Accidental.
Galapagos Distribution: San Cristóbal.
References: Hebard, M. et al. (1920), Linsley, E.G. et al. (1966), Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).
24. *Phoetalia pallida* (Brunner, 1865)
Taxon status: Accepted name; taxon occurs in Galapagos.
 Syn.: *Leurolestes pallidus* (Brunner, 1865)
Origin: Introduced, Accidental.
Galapagos Distribution: San Cristóbal.
References: Hebard, M. et al. (1920), Linsley, E.G. et al. (1966), Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).
25. *Pycnoscelis surinamensis* (Linnaeus, 1758)
Taxon status: Accepted name; taxon occurs in Galapagos.
Origin: Introduced, Accidental.
Galapagos Distribution: Española, Fernandina, Floreana, Isabela, San Cristóbal, Santa Cruz, Santiago.
References: Hebard, M. et al. (1920), Linsley, E.G. et al. (1977), Linsley, E.G. et al. (1966), Peck, S. et al. (1986), Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).
26. *Rhyparobia maderae* (Fabricius, 1781)
Taxon status: Accepted name; taxon occurs in Galapagos.

Origin: Introduced, Accidental.

Galapagos Distribution: Floreana, San Cristóbal.

References: Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).

27. *Symploce pallens* (Stephens, 1835)

Taxon status: Accepted name; taxon occurs in Galapagos.

Syn.: *Symploce lita*, *S. hospes*

Origin: Introduced, Accidental.

Galapagos Distribution: Española, Fernandina, Floreana, Genovesa, Isabela, Marchena, Pinta, San Cristóbal, Santa Cruz, Santa Fé, Santiago.

References: Peck, S.B. et al. (1992), Peck, S.B. et al. (1996), Peck, S.B. et al. (1998), Peck, S.B. et al. (2001).

References:

1. Bahder B.W., Scheffrahn R.H., Kreck J., Keil C., Whitney-King S. (2009) *Termites (Isoptera: Kalotermitidae, Rhinotermitidae, Termitidae) of Ecuador*. Ann. soc. entomol. Fr. (n.s.) 45(4): 529-536.
2. Hebard, M. (1920) *Expedition of the California Academy of Sciences to the Galapagos Islands, 1905-1906 17 Dermaptera and Orthoptera*. Proceedings of the California Academy of Sciences Fourth Series 2(17): 311-346.
3. Light, S.F (1935) *The Templeton Crocker Expedition of the California Academy of Sciences, 1932 No.20 The Termites*. Proceedings of the California Academy of Sciences Fourth Series Vol 21(20): 233-258.
4. Linsley, E.G., Usinger, R.L. (1966) *Insects of the Galápagos Islands*. Proceedings of the California Academy of Sciences Fourth Series 33(7): 113-196.
5. Linsley, E.G. (1977) *Insects of the Galápagos (Supplement)*. Occasional Papers of the California Academy of Sciences 125: 1-50.
6. Peck, S., Peck, J. (1986) *The Galapagos Islands volcanic caves and cave fauna of the Galapagos Islands*. The Canadian Caver 18(1): 42-49.
7. Peck, S.B. (1990) *Eyeless arthropods of the Galapagos Islands, Ecuador: Composition and origin of the cryptozoic fauna of a young, tropical, oceanic archipelago*. Biotropica 22(4): 366-381.
8. Peck, S.B., Roth, L.M. (1992) *Cockroaches of the Galápagos Islands, Ecuador, with descriptions of three new species (Insecta: Blattodea)*. Canadian Journal of Zoology 70: 2202-2217.
9. Peck, S.B. (1996) *Diversity and distribution of the orthopteroid insects of the Galápagos Islands, Ecuador*. Canadian Journal of Zoology 74: 1497-1510.
10. Peck, S.B., Heraty, J., Landry, B. Sinclair, B.J. (1998) *Introduced insect fauna of an oceanic archipelago: The Galápagos Islands, Ecuador*. Am. Entomol. 44: 218-237.
11. Peck, S.B. (2001) *Small Orders of Insects of the Galápagos Islands, Ecuador: Evolution, Ecology, and Diversity*. NRC Research Press, Ottawa, Ontario, Canada, 278 pp.
12. Roth, L.M. (1988) *Some cavernicolous and epigeal cockroaches with six new species, and a discussion of the Nocticolidae (Dictyoptera: Blattaria)*. Revue Suisse Zool. 95(1): 297-321.